

The maximum size of 3-wise intersecting and 3-wise union families

Peter Frankl

CNRS, ER 175 Combinatoire,
2 Place Jussieu, 75005 Paris, France
Peter111F@aol.com

Norihide Tokushige*

College of Education, Ryukyu University,
Nishihara, Okinawa, 903-0213 Japan
hide@edu.u-ryukyu.ac.jp

July 29, 2005

Abstract

Let \mathcal{F} be an n -uniform hypergraph on $2n$ vertices. Suppose that $|F_1 \cap F_2 \cap F_3| \geq 1$ and $|F_1 \cup F_2 \cup F_3| \leq 2n - 1$ holds for all $F_1, F_2, F_3 \in \mathcal{F}$. We prove that the size of \mathcal{F} is at most $\binom{2n-2}{n-1}$.

1 Introduction

A family $\mathcal{F} \subset 2^X$ is called r -wise intersecting if $F_1 \cap \dots \cap F_r \neq \emptyset$ holds for all $F_1, \dots, F_r \in \mathcal{F}$. A family $\mathcal{F} \subset 2^X$ is called r -wise union if $F_1 \cup \dots \cup F_r \neq X$ holds for all $F_1, \dots, F_r \in \mathcal{F}$. The Erdős–Ko–Rado theorem[2] states that if $n \geq 2k$ and $\mathcal{F} \subset \binom{X}{k}$ is 2-wise intersecting then $|\mathcal{F}| \leq \binom{n-1}{k-1}$. By considering

*The second author was supported by MEXT Grant-in-Aid for Scientific Research (B) 16340027.

the complement, the theorem can be restated as follows: if $n \leq 2k$ and $\mathcal{F} \subset \binom{[n]}{k}$ is 2-wise union then $|\mathcal{F}| \leq \binom{n-1}{k}$.

We can extend the Erdős–Ko–Rado theorem for r -wise intersecting families as follows.

Theorem 1 [3] *If $\mathcal{F} \subset \binom{[n]}{k}$ is r -wise intersecting and $(r-1)n \geq rk$ then $|\mathcal{F}| \leq \binom{n-1}{k-1}$. If $r \geq 3$ then equality holds iff $\mathcal{F} = \{F \in \binom{[n]}{k} : i \in F\}$ holds for some $i \in [n]$.*

The equivalent complement version is the following. If $\mathcal{F} \subset \binom{[n]}{k}$ is r -wise union and $rk \geq n$ then $|\mathcal{F}| \leq \binom{n-1}{k}$.

Gronau[6], and Engel and Gronau[1] proved the following.

Theorem 2 *Let $r \geq 4$, $s \geq 4$ and $\mathcal{F} \subset \binom{[n]}{k}$. Suppose that \mathcal{F} is r -wise intersecting and s -wise union, and*

$$\frac{n-1}{s} + 1 \leq k \leq \frac{r-1}{r}(n-1).$$

Then we have $|\mathcal{F}| \leq \binom{n-2}{k-1}$.

In this note we prove the following.

Theorem 3 *Let $\mathcal{F} \subset \binom{[2n]}{n}$ be a 3-wise intersecting and 3-wise union family. Then we have $|\mathcal{F}| \leq \binom{2n-2}{n-1}$. Equality holds iff $\mathcal{F} = \{F \in \binom{[2n]-\{j\}}{n} : i \in F\}$ holds for some $i, j \in [2n]$.*

2 Proof of Theorem 3

We can prove the theorem for $n \leq 3$ easily, so we assume that $n \geq 4$. Let $\mathcal{F} \subset \binom{[2n]}{n}$ be a 3-wise intersecting and 3-wise union family. If $\mathcal{F} \subset \binom{[2n]-\{j\}}{n}$ holds for some $j \in [2n]$ then Theorem 1 implies that $|\mathcal{F}| \leq \binom{2n-2}{n-1}$ and equality holds iff there exists some $i \in [2n]$ such that $i \in F$ holds for all $F \in \mathcal{F}$, which verifies the theorem. From now on we assume that there is no such j , in other words, we assume that

$$\bigcup_{F \in \mathcal{F}} F = [2n]. \tag{1}$$

Considering the complement, we may assume that

$$\bigcap_{F \in \mathcal{F}} F = \emptyset. \quad (2)$$

Now suppose that

$$|\mathcal{F}| \geq \binom{2n-2}{n-1} \quad (3)$$

and we shall prove that there is no such \mathcal{F} .

For $A \in \binom{[2n]}{n}$, we define the corresponding walk on \mathbb{Z}^2 , denoted by $\text{walk}(A)$, in the following way. The walk is from $(0, 0)$ to (n, n) with $2n$ steps, and if $i \in A$ (resp. $i \notin A$) then we move one unit up (resp. one unit to the right) at the i -th step. Let us define

$$\mathcal{A}_i := \{A \in \binom{[2n]}{n} : |A \cap [1 + 3\ell]| \geq 1 + 2\ell \text{ first holds at } \ell = i\},$$

$$\mathcal{A}_{\bar{j}} := \{A \in \binom{[2n]}{n} : |A \cap [2n - 3\ell, 2n]| \leq \ell \text{ first holds at } \ell = j\}.$$

If $A \in \mathcal{A}_i$ then, after starting from the origin, $\text{walk}(A)$ touches the line $y = 2x + 1$ at $(i, 2i + 1)$ for the first time. If $A \in \mathcal{A}_{\bar{j}}$ then $\text{walk}(A)$ touches the line $y = \frac{1}{2}(x - (n - 1)) + n$ at $(n - 2j - 1, n - j)$ and after passing this point this walk never touches the line again. Set $\mathcal{A}_{i\bar{j}} := \mathcal{A}_i \cap \mathcal{A}_{\bar{j}}$, and

$$a_i := |\mathcal{A}_i| / \binom{2n-2}{n-1}, \quad a_{\bar{j}} := |\mathcal{A}_{\bar{j}}| / \binom{2n-2}{n-1}, \quad a_{i\bar{j}} := |\mathcal{A}_{i\bar{j}}| / \binom{2n-2}{n-1}.$$

Set also

$$\begin{aligned} \mathcal{F}_i &:= \mathcal{A}_i \cap \mathcal{F}, & \mathcal{F}_{\bar{j}} &:= \mathcal{A}_{\bar{j}} \cap \mathcal{F}, & \mathcal{F}_{i\bar{j}} &:= \mathcal{A}_{i\bar{j}} \cap \mathcal{F}, \\ f_i &:= |\mathcal{F}_i| / \binom{2n-2}{n-1}, & f_{\bar{j}} &:= |\mathcal{F}_{\bar{j}}| / \binom{2n-2}{n-1}, & f_{i\bar{j}} &:= |\mathcal{F}_{i\bar{j}}| / \binom{2n-2}{n-1}, \end{aligned}$$

and

$$\mathcal{G}_{i\bar{j}} := \{F \cap [3i + 2, 2n - 3j - 1] : F \in \mathcal{F}_{i\bar{j}}\}.$$

Note that $|\mathcal{G}_{i\bar{j}}| \leq |\mathcal{F}_{i\bar{j}}|$ and equality holds if both of i and j are at most 1.

We also use the following basic facts about shifting. (See e.g., [8, 4, 5] for the details.) We may assume that $\mathcal{F} \subset \binom{[2n]}{n}$ is shifted, i.e., for all $F \in \mathcal{F}$ and $1 \leq i < j \leq 2n$, if $i \notin F$ and $j \in F$ then $(F - \{j\}) \cup \{i\} \in \mathcal{F}$. It follows then for all $F \in \mathcal{F}$, $\text{walk}(F)$ must touch the line $y = 2x + 1$ because \mathcal{F} is a shifted 3-wise 1-intersecting family. In the same way, $\text{walk}(F)$ must touch the line $y = \frac{1}{2}(x - (n - 1)) + n$ because \mathcal{F} is a shifted 3-wise 1-union family.

Claim 1 $\mathcal{G}_{0\bar{0}} \subset \binom{[2, 2n-1]}{n-1}$ is 2-wise intersecting.

Proof. Otherwise we have $A, B \in \mathcal{F}_{0\bar{0}}$ such that $A \cap B = \{1\}$. This forces $\bigcap_{F \in \mathcal{F}} F = \{1\}$, contradicting (2). \square

By Claim 1 and the Erdős–Ko–Rado theorem, we have $|\mathcal{F}_{0\bar{0}}| = |\mathcal{G}_{0\bar{0}}| \leq \binom{2n-3}{n-2}$ and

$$f_{0\bar{0}} \leq \binom{2n-3}{n-2} / \binom{2n-2}{n-1} = \frac{1}{2}. \quad (4)$$

Claim 2 $\mathcal{G}_{1\bar{0}} \subset \binom{[5, 2n-1]}{n-3}$ is 2-wise intersecting.

Proof. Suppose on the contrary that there exist $A, B \in \mathcal{G}_{1\bar{0}}$ such that $A \cap B = \emptyset$. Then $\{2, 3, 4\} \cup A, \{2, 3, 4\} \cup B \in \mathcal{F}_{1\bar{0}}$. Since \mathcal{F} is shifted we also have $\{1, 3, 4\} \cup B \in \mathcal{F}_{1\bar{0}}$. If there is $F \in \mathcal{F}$ such that $|F \cap [4]| \leq 2$ then we may assume that $F \cap [2] = \{1, 2\}$ by the shiftedness of \mathcal{F} . But this is impossible because $(\{2, 3, 4\} \cup A) \cap (\{1, 3, 4\} \cup B) \cap F = \emptyset$.

Thus we may assume that $|F \cap [4]| \geq 3$ holds for all $F \in \mathcal{F}$. Let

$$\begin{aligned} \mathcal{F}(\bar{1}234) &:= \{F \cap [5, 2n] : F \in \mathcal{F}, F \cap [4] = \{2, 3, 4\}\} \subset \binom{[5, 2n]}{n-3}, \\ \mathcal{F}(1\bar{2}34) &:= \{F \cap [5, 2n] : F \in \mathcal{F}, F \cap [4] = \{1, 3, 4\}\} \subset \binom{[5, 2n]}{n-3}, \\ \mathcal{F}(12\bar{3}4) &:= \{F \cap [5, 2n] : F \in \mathcal{F}, F \cap [4] = \{1, 2, 4\}\} \subset \binom{[5, 2n]}{n-3}. \end{aligned}$$

Then $|\mathcal{F}(\bar{1}234)| + |\mathcal{F}(1\bar{2}34)| + |\mathcal{F}(12\bar{3}4)| \leq 3 \binom{2n-4}{n-3}$. Let

$$\mathcal{F}(123) := \{F \cap [4, 2n] : \{1, 2, 3\} \subset F \in \mathcal{F}\} \subset \binom{[4, 2n]}{n-3}.$$

Then $\mathcal{F}(123)$ is 3-wise union and it follows from the complement version of Theorem 1 that $|\mathcal{F}(123)| \leq \binom{2n-4}{n-3}$. Therefore we have

$$|\mathcal{F}| = |\mathcal{F}(\bar{1}234)| + |\mathcal{F}(1\bar{2}34)| + |\mathcal{F}(12\bar{3}4)| + |\mathcal{F}(123)| \leq 4 \binom{2n-4}{n-3} < \binom{2n-2}{n-1},$$

which contradicts (3). \square

By Claim 2 and the Erdős–Ko–Rado theorem, we have $|\mathcal{F}_{1\bar{0}}| = |\mathcal{G}_{1\bar{0}}| \leq \binom{2n-6}{n-4}$ and

$$f_{1\bar{0}} \leq \binom{2n-6}{n-4} / \binom{2n-2}{n-1} = \frac{(n-1)(n-3)}{4(2n-3)(2n-5)}.$$

Considering the complement, we have the same estimation for $f_{0\bar{1}}$. Therefore we have

$$f_{1\bar{0}} + f_{0\bar{1}} \leq \frac{(n-1)(n-3)}{2(2n-3)(2n-5)}. \quad (5)$$

Claim 3 $\mathcal{G}_{1\bar{1}} \subset \binom{[5, 2n-4]}{n-4}$ is 2-wise intersecting.

Proof. Suppose that there are $A, B \in \mathcal{G}_{1\bar{1}}$ such that $A \cap B = \emptyset$. Then we have $F_1 := \{2, 3, 4, 2n\} \cup A \in \mathcal{F}$. Since \mathcal{F} is shifted and $\{2, 3, 4, 2n\} \cup B \in \mathcal{F}$, we also have $F_2 := \{1, 3, 4, 2n-1\} \cup B \in \mathcal{F}$. If $|F \cap [4]| \geq 3$ holds for all $F \in \mathcal{F}$ then we are done as we saw in the proof of Claim 2. So there is $G \in \mathcal{F}$ such that $|G \cap [4]| \leq 2$ and by the shiftedness we may assume that $G \cap [4] = \{1, 2\}$. Then $F_1 \cap F_2 \cap G = \emptyset$, which is a contradiction. \square

By Claim 3 and the Erdős–Ko–Rado theorem, we have $|\mathcal{F}_{1\bar{1}}| = |\mathcal{G}_{1\bar{1}}| \leq \binom{2n-9}{n-5}$ and

$$f_{1\bar{1}} \leq \binom{2n-9}{n-5} / \binom{2n-2}{n-1} = \frac{(n-1)(n-2)(n-3)}{16(2n-3)(2n-5)(2n-7)}. \quad (6)$$

By (4), (5) and (6), we have the following.

Claim 4 $f_{0\bar{0}} + f_{1\bar{0}} + f_{0\bar{1}} + f_{1\bar{1}} \leq H_1$, where

$$H_1 := \frac{1}{2} + \frac{(n-1)(n-3)}{2(2n-3)(2n-5)} + \frac{(n-1)(n-2)(n-3)}{16(2n-3)(2n-5)(2n-7)}.$$

Next we consider $f_{i\bar{j}}$ where $\max\{i, j\} = 2$. Let c_i be the number of walks from $(0, 0)$ to $(i, 2i+1)$ which touch the line $y = 2x + 1$ only at $(i, 2i+1)$. Then it follows that $c_i = \frac{1}{3i+1} \binom{3i+1}{i}$ (see e.g. Fact 3 in [7]).

If $A \in \mathcal{A}_{i\bar{j}}$ then walk(A) goes through the two points $P = (i, 2i+1)$ and $Q = (n-2j-1, n-j)$. Since the number of walks from P to Q is $\binom{2n-(3i+3j+2)}{n-(i+2j+1)}$, we get the following simple estimation.

$$f_{i\bar{j}} \leq a_{i\bar{j}} = c_i c_j \binom{2n-(3i+3j+2)}{n-(i+2j+1)} / \binom{2n-2}{n-1} =: g(i, j).$$

Thus we have

$$(f_{2\bar{0}} + f_{0\bar{2}}) + (f_{2\bar{1}} + f_{1\bar{2}}) + f_{2\bar{2}} \leq 2(g(2, 0) + g(2, 1)) + g(2, 2) =: H_2. \quad (7)$$

Finally we consider $f_i, f_{\bar{i}}$ for $i \geq 3$. We use the following fact which we prove in the next section.

Lemma 1 We have

$$\sum_{i=0}^{\lfloor \frac{n-1}{2} \rfloor} |\mathcal{A}_i| \leq \alpha \binom{2n}{n}$$

for all $n \geq 1$ where $\alpha = \frac{\sqrt{5}-1}{2}$.

We also use the following trivial estimation.

$$\max\{f_i, f_{\bar{i}}\} \leq a_i = a_{\bar{i}} = c_i \binom{2n-3i-1}{n-i} / \binom{2n-2}{n-1}.$$

Then this together with Lemma 1 implies

$$\sum_{i>2} f_i \leq \sum_{i>2} a_i \leq \alpha \binom{2n}{n} / \binom{2n-2}{n-1} - \sum_{i=0}^2 a_i =: H_3 \quad (8)$$

By Claim 4, (7) and (8), we have

$$|\mathcal{F}| / \binom{2n-2}{n-1} \leq \sum_{0 \leq i \leq 2, 0 \leq j \leq 2} f_{i\bar{j}} + \sum_{i>2} f_i + \sum_{j>2} f_j \leq H_1 + H_2 + 2H_3 =: H_4(n),$$

where

$$\begin{aligned} H_4(n) = & 4\sqrt{5} - \frac{32551}{4096} - \frac{2(\sqrt{5}-2)}{n} + \frac{1}{2^{20}} \left(\frac{6237}{2n-13} + \frac{2835}{2n-11} + \right. \\ & \left. + \frac{28770}{2n-9} - \frac{156090}{2n-7} + \frac{923313}{2n-5} + \frac{298295}{2n-3} \right). \end{aligned}$$

Note that $\lim_{n \rightarrow \infty} H_4(n) = 4\sqrt{5} - \frac{32551}{4096} = 0.997\dots$. In fact one can check that $H_4(n) < 1$ for $n \geq 34$. For the remainder cases $4 \leq n \leq 33$, one can directly check that

$$|\mathcal{F}| / \binom{2n-2}{n-1} \leq H_1 + H_2 + 2 \sum_{i=3}^{\lfloor \frac{n-1}{2} \rfloor} a_i < 1.$$

Consequently we showed that $|\mathcal{F}| < \binom{2n-2}{n-1}$ for all $n \geq 4$ and this contradicts (3). This completes the proof of Theorem 3. \square

3 Proof of Lemma 1

Since $|\mathcal{A}_i| = c_i \binom{2n-3i-1}{n-i}$ we need to prove that

$$\sum_{i=0}^{\lfloor \frac{n-1}{2} \rfloor} c_i \binom{2n-3i-1}{n-i} / \binom{2n}{n} \leq \alpha.$$

We use the following fact (cf. (6) in [7]):

$$\sum_{i=0}^{\lfloor \frac{n-1}{2} \rfloor} c_i \left(\frac{1}{2}\right)^{3i+1} \leq \sum_{i=0}^{\infty} c_i \left(\frac{1}{2}\right)^{3i+1} = \alpha.$$

Thus to prove the lemma, it suffices to show that

$$\binom{2n-3i-1}{n-i} / \binom{2n}{n} \leq \left(\frac{1}{2}\right)^{3i+1} \quad (9)$$

for $0 \leq i \leq \lfloor \frac{n-1}{2} \rfloor$. We prove this inequality by induction on i . For the case $i = 0$, one can check that the equality holds in (9). Now let $i > 0$ and we assume (9) for i and we show the case $i + 1$, that is,

$$\binom{2n-3i-4}{n-i-1} / \binom{2n}{n} \leq \left(\frac{1}{2}\right)^{3i+4},$$

or equivalently,

$$\binom{2n}{n} \geq 2^{3i+4} \binom{2n-3i-4}{n-i-1}.$$

By the induction hypothesis, we have

$$\binom{2n}{n} \geq 2^{3i+1} \binom{2n-3i-1}{n-i},$$

and so it suffices to show that

$$2^{3i+1} \binom{2n-3i-1}{n-i} \geq 2^{3i+4} \binom{2n-3i-4}{n-i-1},$$

or equivalently,

$$f(i) := 5i^3 - (10n+6)i^2 + (4n^2-17)i + 6n - 6 \geq 0.$$

Since $f''(i) = -2(10n - 15i + 6) < 0$, the function $f(i)$ is concave on the domain $0 \leq i \leq \lfloor \frac{n-1}{2} \rfloor$. Thus it suffices to check that $f(0) \geq 0$ and $f(\lfloor \frac{n-1}{2} \rfloor) \geq 0$. Indeed, $f(0) = 6(n-1) \geq 0$, and $f(\lfloor \frac{n-1}{2} \rfloor) \geq \min\{f(\frac{n-1}{2}), f(\frac{n-2}{2})\} = f(\frac{n-1}{2}) = \frac{1}{8}(n+1)(n-1)(n-3) \geq 0$ if $n \geq 3$. For the case $n \leq 2$, we only have $0 \leq i \leq \lfloor \frac{1}{2} \rfloor = 0$, that is, $i = 0$ and we already checked this case. \square

Acknowledgment. The authors would like to thank Professor Konrad Engel for telling them the problem and related references.

References

- [1] K. Engel, H.-D.O.F. Gronau. An Erdős–Ko–Rado type theorem II. *Acta Cybernet.*, 4:405–411, 1986.
- [2] P. Erdős, C. Ko, R. Rado. Intersection theorems for systems of finite sets. *Quart. J. Math. Oxford (2)*, 12:313–320, 1961.
- [3] P. Frankl. On Sperner families satisfying an additional condition. *J. Combin. Theory (A)*, 20:1–11, 1976.
- [4] P. Frankl, N. Tokushige. Weighted 3-wise 2-intersecting families. *J. Combin. Theory (A)*, 100:94–115, 2002.
- [5] P. Frankl, N. Tokushige. Random walks and multiply intersecting families. *J. Combin. Theory (A)*, 109:121–134, 2005.
- [6] H.-D.O.F. Gronau. An Erdős–Ko–Rado type theorem. Finite and infinite sets, Vol. I,II (Eger, 1981) *Colloq. Math. Soc. J. Bolyai*, 37:333–342, 1984.
- [7] N. Tokushige. A frog’s random jump and the Pólya identity. *Ryukyu Math. Journal*, 17:89–103, 2004.
- [8] N. Tokushige. The maximum size of 4-wise 2-intersecting and 4-wise 2-union families. *European J. Combin.*, in press.